

“QCD AND STANDARD MODEL”
Problem Set 7

Non-Abelian gauge theories

Non-Abelian gauge theories play an important role in particle physics. In this exercise we shall discuss some properties of non-Abelian gauge theories with group $SU(N)$.

- a) The Yang-Mills Lagrangian density for a non-Abelian gauge field A_μ^a can be written in the following form,

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4}F_{\mu\nu}^a F^{\mu\nu a},$$

with $F_{\mu\nu}^a \equiv \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc}A_\mu^b A_\nu^c$, f^{abc} the structure constants of $SU(N)$ and summation over all repeated indexes is tacitly assumed. Show that the above can be equivalently written as

$$\mathcal{L}_{\text{YM}} = -\frac{1}{2}\text{Tr}(F_{\mu\nu}F^{\mu\nu}),$$

where $F_{\mu\nu} \equiv F_{\mu\nu}^a T^a$, with T^a the generators of $SU(N)$.

- b) Consider now

$$\mathcal{L} = \mathcal{L}_{\text{YM}} + \mathcal{L}_{\text{D}},$$

where the generalized Dirac Lagrangian reads

$$\mathcal{L}_{\text{D}} = \bar{\psi}(i\gamma^\mu D_\mu)\psi - m\bar{\psi}\psi,$$

with the covariant derivative $D_\mu \equiv \partial_\mu - igA_\mu$, and $A_\mu \equiv A_\mu^a T^a$. Show that \mathcal{L} is invariant under the Yang-Mills transformation

$$\begin{aligned} A_\mu(x) &\longrightarrow A'_\mu(x) = U(x)A_\mu(x)U^{-1}(x) - \frac{i}{g}[\partial_\mu U(x)]U^{-1}(x), \\ \psi(x) &\longrightarrow \psi'(x) = U(x)\psi(x), \end{aligned}$$

for any $SU(N)$ matrix $U(x)$.

- c) Take a global $SU(N)$ transformation, i.e. $U = \text{const.}$ and find the conserved current using the Noether theorem. Compare the corresponding charge to the one you found in the Abelian case in problem set 1.
- d) Derive the equations of motion for A_μ^a following from \mathcal{L} and compare them with the corresponding equations of motion for an Abelian gauge field A_μ in QED.
- e) Show that

$$[D_\mu, D_\nu]\psi = -igF_{\mu\nu}\psi,$$

where the brackets $[\dots]$ stand for the commutator.

- f) Demonstrate the *Bianchi identity*

$$D_\rho F_{\mu\nu} + D_\mu F_{\nu\rho} + D_\nu F_{\rho\mu} = 0.$$

- g) Consider now the dual field strength tensor defined as

$$\tilde{F}_{\mu\nu} \equiv \frac{1}{2}\epsilon_{\mu\nu\alpha\beta}F^{\alpha\beta}.$$

First, show that in the Abelian case

$$F^{\mu\nu} \tilde{F}_{\mu\nu} = \partial_\mu K^\mu,$$

and find K_μ . Does this term have any consequences for the equations of motion?

Next, show that the corresponding quantity in the non-Abelian case (which has to be SU(N)-invariant) is

$$\text{Tr} \left(F^{\mu\nu} \tilde{F}_{\mu\nu} \right) = \partial_\mu \tilde{K}^\mu.$$

What is \tilde{K}_μ now?

Discuss whether a term like $\theta \text{Tr} \left(F^{\mu\nu} \tilde{F}_{\mu\nu} \right)$, with θ a constant, is allowed in the Lagrangian. Does it appear in the equations of motion? What are the implications?