

## Sheet 3:

Hand-out: Friday, May 13, 2022<sup>1</sup>

### Problem 1 Bogoliubov theory of the Bose-Hubbard model

In this problem, we extend the Bogoliubov theory of weakly interacting bosons to a BEC in a lattice model. You can follow closely the continuum calculation from the lecture. Specifically, consider the Bose-Hubbard Hamiltonian in 3D,

$$\hat{\mathcal{H}} = -t \sum_{\langle i,j \rangle} (\hat{a}_i^\dagger \hat{a}_j + \text{h.c.}) + \frac{U}{2} \sum_j (\hat{a}_j^\dagger \hat{a}_j - 1) \hat{a}_j^\dagger \hat{a}_j, \quad (1)$$

where  $[\hat{a}_i, \hat{a}_j^\dagger] = \delta_{i,j}$  are bosonic operators.

- (1.a) Determine the normal-ordered Hamiltonian :  $\hat{\mathcal{H}}$  : and show that :  $\hat{\mathcal{H}} := \hat{\mathcal{H}}$ . Explain the physical meaning of the different terms in the Hamiltonian.
- (1.b) Introduce discrete momentum modes, associated with second quantized bosonic operators  $\hat{a}_{\mathbf{k}}$ , and make a variational ansatz which describes a macroscopic occupation of the  $\mathbf{k} = 0$  mode.
- (1.c) Derive the effective Hamiltonian, quadratic in  $\hat{a}_{\mathbf{k} \neq 0}$ , describing low-energy collective excitations.
- (1.d) Diagonalize the effective Hamiltonian from (1.c) and derive the Bogoliubov dispersion relation in a lattice.

### Problem 2 Classical field theory

In this problem we discuss *classical field theories* and their formulation using Lagrangian densities. Later in the lecture we focus on *quantum field theories*, but it is often useful to relate them (in some limits) to simpler classical theories.

A classical field  $\phi_n(\mathbf{r}, t)$  with components  $n$  is described with a general action of the form:

$$S = \int dt \int d^d \mathbf{r} \mathcal{L}[\phi_n, \partial_\mu \phi_n], \quad (2)$$

where  $\mu = t, x, y, z, \dots$  and  $\mathcal{L}[\phi_n, \partial_\mu \phi_n]$  is the Lagrangian density. By minimizing the action,  $\delta S = 0$ , one obtains the Euler-Lagrange equations of motion:

$$\sum_\mu \frac{\partial}{\partial x_\mu} \frac{\partial \mathcal{L}[\phi_n, \partial_\mu \phi_n]}{\partial (\partial_\mu \phi_n)} - \frac{\partial \mathcal{L}[\phi_n, \partial_\mu \phi_n]}{\partial \phi_n} = 0 \quad \forall n. \quad (3)$$

<sup>1</sup>If you would like to present your solution(s), feel free to send them to Henning Schlömer until Fri, May 20.

- (2.a) Consider the classical  $\phi^4$  theory of a *real, scalar, non-relativistic* field  $\phi \in \mathbb{R}$  described by the action

$$\mathcal{L}[\phi, \partial_\mu \phi] = \frac{1}{2} \sum_\mu (\partial_\mu \phi)^2 - \frac{m^2}{2} \phi^2 - \alpha \phi^4, \quad (4)$$

and derive the corresponding equations of motion for  $\phi(\mathbf{r}, t)$ .

- (2.b) Consider the classical  $\phi^4$  theory of a *complex, scalar, non-relativistic* field  $\phi \in \mathbb{C}$  described by the action

$$\mathcal{L}[\phi, \phi^*, \partial_\mu \phi, \partial_\mu \phi^*] = \frac{1}{2} \sum_\mu |\partial_\mu \phi|^2 - \frac{m^2}{2} |\phi|^2 - \alpha |\phi|^4, \quad (5)$$

and derive the corresponding equations of motion for  $\phi(\mathbf{r}, t)$  and  $\phi^*(\mathbf{r}, t)$ . You may treat  $\phi$  and  $\phi^*$  as two independent components!

### Problem 3 The Gross-Pitaevskii equation

In this problem we study weakly interacting bosons with point-like interactions of strength  $g$  in an external trapping potential  $V(\mathbf{x})$ . With  $m$  and  $\mu$  we denote the boson mass and chemical potential, respectively. This system is described by the Hamiltonian ( $\hbar = 1$ ):

$$\hat{\mathcal{H}} = \int d^3\mathbf{r} \left\{ \frac{1}{2m} |\nabla \hat{\psi}(\mathbf{r})|^2 + (V(\mathbf{r}) - \mu) \hat{\psi}^\dagger(\mathbf{r}) \hat{\psi}(\mathbf{r}) \right\} + \frac{g}{2} \int d^3\mathbf{r} d^3\mathbf{r}' \hat{\psi}^\dagger(\mathbf{r}) \hat{\psi}^\dagger(\mathbf{r}') \delta(\mathbf{r} - \mathbf{r}') \hat{\psi}(\mathbf{r}') \hat{\psi}(\mathbf{r}) \quad (6)$$

- (3.a) Using the canonical commutation relations of the bosonic field  $\hat{\psi}(\mathbf{r})$ , derive the equations of motion of the field operators  $\hat{\psi}(\mathbf{r}, t)$  in the Heisenberg picture:

$$i \frac{\partial}{\partial t} \hat{\psi}(\mathbf{r}, t) = [\hat{\psi}(\mathbf{r}, t), \hat{\mathcal{H}}] = \dots \quad (7)$$

You obtain the operator-valued Gross-Pitaevskii equation.

- (3.b) To derive a simpler  $\mathbb{C}$ -valued classical equation providing an approximate description of the interacting Bose-gas, consider the following variational wavefunction:

$$|\psi(t)\rangle = \frac{1}{\sqrt{N!}} \left( \int d^3\mathbf{r} \Psi(\mathbf{r}, t) \hat{\psi}^\dagger(\mathbf{r}) \right)^N |0\rangle. \quad (8)$$

Here  $N$  is the total boson number and  $\Psi(\mathbf{r}, t)$  is a variational parameter depending on space and time. Describe the physical meaning of this state!

- (3.c) For the variational ansatz in (3.b), derive the following expectation value,

$$\mathcal{L}[\partial_t \Psi, \nabla \Psi, \Psi] = \langle \psi(t) | -i\partial_t + \hat{\mathcal{H}} | \psi(t) \rangle \quad (9)$$

which takes the role of a classical Lagrangian density.

- (3.d) Derive the Euler-Lagrange equations for  $\Psi(\mathbf{r}, t)$  from the Lagrangian density  $\mathcal{L}$  derived in (3.c). Show that the obtained equation takes the form:

$$i\partial_t \Psi(\mathbf{r}, t) = -\frac{1}{2m} \nabla^2 \Psi(\mathbf{r}, t) + (V(\mathbf{r}) - \mu) \Psi(\mathbf{r}, t) + g |\Psi(\mathbf{r}, t)|^2 \Psi(\mathbf{r}, t). \quad (10)$$

Compare this *Gross-Pitaevskii equation* to the operator-valued equation of motion obtained in (3.a)!

*Hint:* You may treat space as discretize to conceptually simplify your calculations.