

Ludwig-Maximilians-Universität München

Quantum Field Theory (Quantum Electrodynamics)

Prof. Dr. Georgi Dvali

Assistants : Lukas Eisemann, Oleg Kaikov,
Dr. Georgios Karananas, Max Warkentin

19 February 2021

Guidelines :

- The exam consists of 6 problems.
- The duration of the exam is 24 hours.
- Please write your name or matriculation number on every sheet that you hand in.
- Your answers should be comprehensible and readable.

GOOD LUCK!

Exercise 1	25 P
Exercise 2	20 P
Exercise 3	15 P
Exercise 4	15 P
Exercise 5	20 P
Exercise 6	5 P

Total	100 P
-------	-------

Problem 1 (25 points)

Take the following Lagrangian density in 4 spacetime dimensions (we use units $\hbar = c = 1$),

$$\mathcal{L} = i\bar{\psi}\gamma^\mu\partial_\mu\psi + \lambda((\bar{\psi}\psi)^2 - (\bar{\psi}\gamma_5\psi)^2), \quad (1)$$

where ψ is a spinor and λ is a constant. As usual, $\bar{\psi} \equiv \psi^\dagger\gamma_0$.

- What is the mass dimension of ψ ?
- What is the mass dimension of λ ?
- Find the equations of motion for the theory.
- Requiring that the action is Lorentz invariant, can ψ be a Dirac spinor (in terms of degrees of freedom)?
- Requiring that the action is Lorentz invariant, can ψ be a Majorana spinor?
- Introduce left-handed ψ_L and right-handed ψ_R chiral spinors. How do they transform under $\psi \rightarrow \psi' = e^{i\alpha\gamma_5}\psi$, with α a nonzero real constant?
- Write the Lagrangian density in terms of the ψ_L and ψ_R spinors. Is it invariant under the above chiral transformation? If yes, find the corresponding Noether current. Check that it is conserved on the equations of motion.
- Consider the $\psi\psi \rightarrow \psi\psi$ scattering process. Derive the spin-averaged amplitude squared at the leading order in λ .

Problem 2 (20 points)

Consider the QED Lagrangian density

$$\mathcal{L} = i\bar{\psi}\gamma^\mu D_\mu\psi - m\bar{\psi}\psi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}, \quad (2)$$

where

$$D_\mu \equiv \partial_\mu - ieA_\mu, \quad F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (3)$$

- What is the gauge redundancy of this Lagrangian?
- How is this redundancy affected if we deform the theory by adding a mass term for the vector?

$$\mathcal{L} = i\bar{\psi}\gamma^\mu D_\mu\psi - m\bar{\psi}\psi + \frac{1}{2}m_A^2\tilde{A}_\mu\tilde{A}^\mu - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (4)$$

Notice that a new notation is introduced in order to distinguish between the massive Proca field \tilde{A}_μ from the massless Maxwell field A_μ .

- How many degrees of freedom are propagated by A_μ and \tilde{A}_μ ? Explain.
- Can the massive theory be written in manifestly gauge redundant (gauge invariant) form? If yes, write it.
- Write down the Feynman rule for the vertex of the theory in equation (4).
- What are the possible polarization states of the massive Proca field?

Problem 3 (15 points)

Consider a theory with two real scalar fields, ϕ and χ , with the following Lagrangian density

$$\mathcal{L} = \frac{3}{2}\partial_\mu\phi\partial^\mu\phi + \frac{3}{2}\partial_\mu\chi\partial^\mu\chi + \partial_\mu\phi\partial^\mu\chi - m^2(\phi^2 + \chi^2). \quad (5)$$

- Find the Lagrangian density for canonically normalized fields.
- Quantize this theory and write down the canonical commutation relations.
- Express the Hamiltonian in terms of creation and annihilation operators.

Problem 4 (15 points)

For two spinors, ψ and χ , consider the following quantities

$$\begin{aligned} \text{A)} & \bar{\psi}_L\chi_R \\ \text{B)} & \bar{\psi}_L\gamma_\mu\chi_R \\ \text{C)} & \bar{\psi}_L\gamma_\mu\partial^\mu\chi_R \\ \text{D)} & \bar{\psi}_L\gamma_\mu\partial^\mu\chi_L \\ \text{E)} & \bar{\psi}_L\chi_L \\ \text{F)} & (\bar{\psi}_L\gamma_\mu\chi_L)(\bar{\psi}_L\gamma^\mu\chi_L) \\ \text{G)} & (\bar{\psi}_L\gamma_\mu\chi_L)(\bar{\psi}_R\gamma^\mu\chi_R) \end{aligned} \quad (6)$$

where L, R denote the chiralities (corresponding to the ± 1 eigenvalues of the γ_5 matrix) and for any spinor X , $\bar{X} \equiv X^\dagger\gamma_0$.

- Which of the above quantities are identically zero?
- Which of them can be non-zero and Lorentz-scalars?
- Which of them can be non-zero and Lorentz-vectors?

Justify your answer in each case.

Problem 5 (20 points)

Consider the following Lagrangian density in $d = 4$ spacetime dimensions (we use units $\hbar = c = 1$)

$$\mathcal{L} = \frac{1}{2}\partial_\mu\Phi\partial^\mu\Phi + \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{1}{2}M^2\Phi^2 - \frac{1}{2}m^2\phi^2 - \kappa\Phi\phi^2, \quad (7)$$

which involves two scalar fields Φ and ϕ with masses M and m , respectively, and κ is a constant.

- a) What is the mass dimension of κ ?
- b) What are the conditions on the masses of the particles such that a particle of type Φ can decay into two particles of type ϕ ?
- c) Write down the Feynman rules for this theory.
- d) Consider the decay $\Phi \rightarrow \phi\phi$. Draw the Feynman diagram(s) contributing to this process at tree-level (lowest order). Use it to derive the expression for the amplitude squared.
- e) Compute the lifetime of the particle Φ to lowest order in κ .

Problem 6 (5 points)

Under what circumstances can a massless particle decay into two other massless particles? Explain.